



ELSEVIER

Aquaculture 216 (2003) 391–393

**Aquaculture**

www.elsevier.com/locate/aqua-online

## Contents of *Aquaculture*, Volume 216

VOL. 216 NOS. 1–4

10 February 2003

### Diseases, Parasites and Predators

- Primary cultures of hemocytes from *Mytilus galloprovincialis* Lmk.: expression of IL-2R $\alpha$  subunit  
A. Cao (Lugo, Spain), L. Mercado (Chile), J.I. Ramos-Martinez and R. Barcia (Lugo, Spain). . . . . 1
- Qualitative and quantitative studies on the relative virus load of tails and heads of shrimp acutely infected with WSSV  
S.V. Durand (Tucson, AZ, USA; Montpellier, France), R.M. Redman, L.L. Mohny, K. Tang-Nelson (Tucson, AZ, USA), J.R. Bonami (Montpellier, France) and D.V. Lightner (Tucson, AZ, USA) . . . . . 9
- Induced resistance to white spot syndrome virus infection in *Penaeus stylirostris* through pre-infection with infectious hypodermal and hematopoietic necrosis virus—a preliminary study  
K.F.J. Tang, S.V. Durand, B.L. White, R.M. Redman, L.L. Mohny and D.V. Lightner (Tucson, AZ, USA). . . . . 19
- Effect of fasting on feed intake, growth and mortality of chinook salmon, *Oncorhynchus tshawytscha*, during an induced *Aeromonas salmonicida* epizootic  
J. Pirhonen, C.B. Schreck (Corvallis, OR, USA), P.W. Reno (Newport, OR, USA) and H. Ögüt (Trabzon, Turkey) . . . . . 31

### Genetics and Breeding

- Effects of heat-shock on cell division and microtubule organization in zygotes of the shrimp *Penaeus indicus* (Crustacea, Decapoda) observed with confocal microscopy  
M. Morelli (Nouméa, New Caledonia), Aquacop (Taravao, French Polynesia). . . . . 39
- Production of a high percentage of male offspring with a natural androgen, 11 $\beta$ -hydroxyandrostenedione (11 $\beta$ OHA4), in Florida red tilapia  
D. Desprez (Montpellier, France; Belgium), E. G  raz (Montpellier, France), M.C. Hoareau (France), C. M  lard (Belgium), P. Bosc (France) and J.F. Baroiller (Montpellier, Rennes, France). . . . . 55
- A combined salt and confinement stress enhances mortality in rainbow trout (*Oncorhynchus mykiss*) selected for high stress responsiveness  
S.-E. Fevolden (Troms  , Norway), K.H. R  ed (Oslo, Norway) and K. Fjalestad (Norway) . . . . . 67

### Husbandry and Management

- Identification and growth rates comparison of divergent migratory contingents of Japanese eel (*Anguilla japonica*)  
W.-N. Tzeng, Y. Iizuka, J.-C. Shiao (Taipei, Taiwan), Y. Yamada and H.P. Oka (Japan) . . . . . 77
- A beneficial coculture: charm abalone *Haliotis discus hannai* and sea cucumber *Stichopus japonicus*  
K.H. Kang, J.Y. Kwon and Y.M. Kim (Yosu, South Korea) . . . . . 87
- Transport of silver catfish (*Rhamdia quelen*) fingerlings at different times, load densities, and temperatures  
J.I. Golombieski, L.V.F. Silva, B. Baldisserotto and J.H.S. da Silva (Santa Maria, Brazil). . . . . 95

Do stocking density and feed reward level affect growth and feeding of self-fed juvenile European sea bass?	
M. Paspatis (Heraklion, Greece), T. Boujard (France), D. Maragoudaki (Heraklion, Greece), G. Blanchard (Namur, Belgium) and M. Kentouri (Heraklion, Greece) . . . . .	103
Variability in mussel settlement on suspended ropes placed at Ahipara Bay, Northland, New Zealand	
A.C. Alfaro and A.G. Jeffs (Auckland, New Zealand) . . . . .	115
Growth and survival of the tehuelche scallop <i>Aequipecten tehuelchus</i> in culture	
M.A. Narvarte (Argentina) . . . . .	127
Bioencapsulation in <i>Artemia</i> : II. Influences of the particle concentration in the enrichment process	
R. Gelabert F. (Ciudad del Carmen, Mexico) . . . . .	143
KCl induced paralysis facilitates detachment of hatchery reared juvenile green sea urchins, <i>Strongylocentrotus droebachiensis</i>	
N.T. Hagen (Bodø, Norway). . . . .	155
Food intake of <i>Macrobrachium rosenbergii</i> during larval development	
H.P. de Barros and W.C. Valenti (São Paulo, Brazil) . . . . .	165
<b>Nutrition</b>	
The resistance to physical stresses by <i>Penaeus monodon</i> juveniles fed diets supplemented with astaxanthin	
Y.-H. Chien, C.-H. Pan (Keelung, Taiwan) and B. Hunter (Bangkok, Thailand). . . . .	177
Use of the bacteria <i>Streptococcus faecium</i> and <i>Lactobacillus acidophilus</i> , and the yeast <i>Saccharomyces cerevisiae</i> as growth promoters in Nile tilapia ( <i>Oreochromis niloticus</i> )	
M. Lara-Flores, M.A. Olvera-Novoa, B.E. Guzmán-Méndez and W. López-Madrid (Mérida, Mexico) . . . . .	193
Optimisation of T-ISO biomass production rich in essential fatty acids: I. Effect of different light regimes on growth and biomass production	
I. Tzovenis, N. De Pauw and P. Sorgeloos (Ghent, Belgium) . . . . .	203
Optimisation of T-ISO biomass production rich in essential fatty acids: II. Effect of different light regimes on the production of fatty acids	
I. Tzovenis, N. De Pauw and P. Sorgeloos (Ghent, Belgium) . . . . .	223
Optimum daily ration for Eurasian perch <i>Perca fluviatilis</i> L. reared at its optimum growing temperature	
E.D. Fiogbé (Cotonou, Benin) and P. Kestemont (Namur, Belgium). . . . .	243
Apparent digestibility of selected feedstuffs by mud crab, <i>Scylla serrata</i>	
M.R. Catacutan, P.S. Eusebio (Tigbauan, Philippines) and S.-i. Teshima (Kagoshima City, Japan) . . . . .	253
Determination of fat content in fish feed by supercritical fluid extraction and subsequent lipid classification of extract by thin layer chromatography-flame ionization detection	
R.B. Johnson and H.J. Barnett (Seattle, WA, USA) . . . . .	263
Potato protein concentrate with low content of solanidine glycoalkaloids in diets for Atlantic salmon ( <i>Salmo salar</i> )	
S. Refstie (Sunndalsøra, Norway) and H.A.J. Tiekstra (Veendam, The Netherlands). . . . .	283
Lipid and fatty acid composition of ovaries from wild fish and ovaries and eggs from captive fish of white sea bream ( <i>Diplodus sargus</i> )	
J.R. Cejas, E. Almansa, J.E. Villamandos, P. Badía, A. Bolaños and A. Lorenzo (Santa Cruz de Tenerife, Spain) . . . . .	299
<b>Physiology and Endocrinology</b>	
Gonadal differentiation and effects of temperature on sex determination in southern flounder ( <i>Paralichthys lethostigma</i> )	
J.A. Luckenbach, J. Godwin, H.V. Daniels and R.J. Borski (Raleigh, NC, USA). . . . .	315
Expression of common carp growth hormone in the yeast <i>Pichia pastoris</i> and growth stimulation of juvenile tilapia ( <i>Oreochromis niloticus</i> )	
Y. Li, J. Bai, Q. Jian, X. Ye, H. Lao, X. Li, J. Luo and X. Liang (Guangzhou, PR China) . . . . .	329
Influences of light intensity and photoperiod on moulting and growth of <i>Penaeus merguensis</i> cultured under laboratory conditions	
T. Hoang, M. Barchiesi, S.Y. Lee, C.P. Keenan and G.E. Marsden (Grafton, Australia). . . . .	343

Differences in the biochemical content of buoyant and non-buoyant eggs of the Japanese eel, <i>Anguilla japonica</i> M. Seoka, S. Yamada, Y. Iwata, T. Yanagisawa, T. Nakagawa and H. Kumai (Wakayama, Japan) . .	355
Brain activity of Atlantic salmon ( <i>Salmo salar</i> ) following electrical stunning using various field strengths and pulse durations D.H.F. Robb (Bristol, UK) and B. Roth (Bergen, Norway) . . . . .	363
Induction of sperm activation in open and closed thelycum penaeoid shrimps J. Alfaro, N. Muñoz (Puntarenas, Costa Rica), M. Vargas (San José, Costa Rica) and J. Komen (Wageningen, Netherlands) . . . . .	371
<b>Book review</b>	
The biology of decapod crustacean larvae (Crustacean Issues 14) M.B. New (Marlow, UK) . . . . .	383
<b>Erratum</b>	
Erratum to "Genetic variation for growth rate, feed conversion efficiency, and disease resistance exists within a farmed population of rainbow trout" [Aquaculture 209 (2002) 59–76] M. Henryon, A. Jokumsen, P. Berg, I. Lund, P.B. Pedersen, N.J. Olesen and W.J. Sliereendrecht (Aarhus, Denmark) . . . . .	389
<b>Contents of Aquaculture, Volume 216</b> . . . . .	391

